



reddish, and have surfaces coated with gray and scaly pubescence, becoming smooth.

Once thought to be a beneficial windbreak tree, it since has been deemed detrimental to the environment. Russian olive can grow in a variety of soil and moisture conditions, but prefers open, moist, riparian zones. It is shade tolerant and can be found along streams, floodplains, fields and open areas up to approximately 8,000 feet in elevation. Russian-olive can outcompete native plants, interfere with natural plant succession and nutrient cycling, and tax water reserves. Because Russian olive is capable of fixing nitrogen in its roots, it can grow on bare, mineral substrates and dominate riparian vegetation. Although Russian olive provides a plentiful source of edible fruits for birds, ecologists have found that bird species richness is actually higher in riparian areas dominated by native vegetation.

The key to effective control of Russian olive is preventing establishment of the trees or shrubs. If plants are already present, control options include cut-stump treatments and mechanical mowing. These treatments depend on size and location of the plant. Details on the back of this sheet can help you create a management plan compatible with your site ecology.



Russian olive (*Elaeagnus angustifolia*) is a perennial tree or shrub that is native in Europe and Asia. The plant has olive-shaped fruits, silver color at first then becoming yellow-red when mature. Russian olive can reproduce by seed or root suckers. Seeds are readily spread by birds and can remain viable for up to 3 years. Spring moisture and slightly alkaline soil tend to favor seedling growth. The plant's extensive root system sprouts root suckers frequently. The tree can reach up to 30 feet in height with branches that have 1 to 2 inch thorns. Leaves are 2 to 3 inches long, alternate, narrow, and have simple blades with smooth edges. The leaf's lower surface is silvery white, while the upper surface is light green in color. Flowers are 4 small sepals in light yellow clusters, fragrant, and appear May through June. Fruits mature from September to November. Russian olive twigs are flexible,


# Russian Olive

*Elaeagnus angustifolia*

Russian Olive  
*Elaeagnus angustifolia*

2013 Quarterquad Survey  
Distribution and Abundance  
In Colorado

64,150+ Infested Acres



Distribution Legend: 0 acres 1-10 acres 11-50 acres 51-300 acres 301-999 acres >1000 acres Not Reported  
Acreage estimates supplied by County Weed Coordinators and compiled by the Colorado Department of Agriculture.

Russian olive is redesignated as a “List B” species in the Colorado Noxious Weed Act. It is required to be either eradicated, contained, or suppressed depending on the local infestations. For more information visit [www.colorado.gov/ag/weeds](http://www.colorado.gov/ag/weeds) and click on the Noxious Weed Management Program. Or call the State Weed Coordinator at the Colorado Department of Agriculture, Conservation Services Division, 303-239-4100.

### Key ID Points

1. Leaves are silvery white.
2. Branches have 1 to 2 inch thorns.
3. Yellow-red fruits on mature plants.
4. Mature trees have shedding, reddish-brown bark.

# Integrated Weed Management Recommendations

Integrated weed management offers the most effective combination of control efforts through the “cut stump” treatment. Trees are cut down with a hatchet or chainsaw, then immediately treated with an approved herbicide on the surface of the cut stump. The most effective timing is late summer/early fall for herbicide transfer into the roots.



© John Randall, TNC



© James Miller, USFS



© Chris Ness, Adams County



© Scott Peterson, USDA

## Russian olive *Elaeagnus angustifolia*

### CULTURAL

Replace Russian olives with native trees. Prevent establishment of new trees by removing seedlings and saplings before they mature. Contact your local Natural Resources Conservation Service for recommendations of other possible trees or shrubs.

### BIOLOGICAL

Tubercularia canker is an unapproved biocontrol. However, it overwinters on infected stems and spreads via rain-splash, animals, or pruning implements to open wounds in the bark. Infected tissue becomes discolored or sunken. Entire stems may be girdled and killed, and the disease can deform or kill stressed plants over time.

### MECHANICAL

Saplings can be pulled with a weed-wrench or cut with brush-cutters. Trees can be girdled or cut with chainsaws. However, stump sprouting commonly occurs after cutting down the tree; and stump excavation without removing all parts of the roots can result in root sprouting. Treating cut-stumps with an herbicide can eliminate sprouting. Stump burning is practical when conditions support a long, hot fire and most effective in summer or early fall. Saplings are most sensitive to mechanical treatment.

### CHEMICAL

The table below includes recommendations for herbicides that can be applied to range and pasturelands. Always read, understand, and follow the label directions. The herbicide label is the LAW!

Herbicide	Rate	Application Timing
Triclopyr (Garlon 4, Remedy)	20-30% solution in basal bark oil. The herbicide Pathfinder and does not require dilution.	Cut-Stump Treatment: Apply to the cambial layer of the tree immediately after the cut-stump treatment and to roots above soil surface. (Summer to fall; fall treatments showed fewer re-growth) Basal Bark Treatment: Spray till wet but not dripping; the roots above soil surface, root collar, and lower trunk to a height of 12-15 inches above ground (Late summer to fall)
Glyphosate* (Rodeo - approved aquatic label)	Undiluted (100% solution) or 50% solution in basal bark oil	Cut-Stump Treatment: Apply to the cambial layer of the tree immediately after the cut-stump treatment and to roots above soil surface. Diluted solutions requires regular agitation. Treat summer to fall; fall treatments showed fewer re-growth.

Note: \*These products are non-selective and will kill any vegetation contacted.

Additional herbicide recommendations for this and other species can be found at:  
[www.colorado.gov/agconservation/CSUHerbicideRecommendations.pdf](http://www.colorado.gov/agconservation/CSUHerbicideRecommendations.pdf)